

## Muscle Contraction

## Control of Body Movements

Muscle groups occur in pairs which work both against and with each other. These opposing groups are located on the opposite side of a joint and perform opposite actions on it.

Agonist: (protagonist) responsible for the primary joint motion  
Antagonist: acts against the agonist to return the joint to extended position

As one group is in full contraction, the other group is in full extension

Example: Bicep Curl

Biceps (agonist) vs Triceps (antagonist)

Agonist and Antagonist groups work together for fine motor control and balance of movement

Sep 24-12:10 PM

## Assisting Muscle Groups

Synergist: allows primary movement of the joint by helping the agonist

*Brachioradialis*

Fixator: fixates the origin from moving

*Laticinus Dors.*

## Reciprocal Innervation

The innervation process which produces exactly the amount of relaxation in the antagonist to balance the amount of contraction generated in the agonist is known as reciprocal innervation. Failure of the central nervous system (CNS) to coordinate the nerve actions of the agonist and antagonist will result in jerky, imprecise movements lacking the smoothness we take for granted.

## Parkinson. Disease

Sep 24-1:13 PM

## 3 Types of Muscle Contraction (Activation)

Concentric Contraction: muscle length decreases against resistance (ie. weight or force ie. gravity)

Eccentric Contraction: muscle length increases against resistance (ie. against weight or force of opposing muscle group)  
(controlled lowering)

Isometric Contraction: muscle length does not change during contraction against a force or weight

Sep 24-1:50 PM

## Muscle Direction: Shortening and Lengthening

Muscle Contraction can only take place in the direction of the muscle fibres:

ie. Abdominals

Rectus Abdominis

Transverse Abdominis

Internal Obliques

External Obliques



Sep 24-1:57 PM